



Rewarding Learning

**General Certificate of Secondary Education
January 2021**

GCSE Mathematics

FOUNDATION TIER ADDITIONAL SUPPORT MATERIALS (For use in January 2021)

FOUNDATION TIER ADDITIONAL SUPPORT MATERIALS (January 2021)

Conversion from imperial to metric units

5 miles = 8 kilometres


Range

The range of a set of data is the difference between the largest value and the smallest value in the data set.

Mean

The mean of a set of data is the sum of all the data values divided by the number of data values.

Perimeter, Area and Volume



The perimeter of a rectangle is the distance around the outside of the rectangle. It is found by adding the lengths of the 4 sides of the rectangle.

$P = 2L + 2B$ where P is perimeter, L is length and B is breadth.

The area of a rectangle is found by multiplying the length of the rectangle by the breadth.

$A = L \times B$ where A is area, L is length and B is breadth.

The volume of a cuboid is found by multiplying the length by the breadth by the height of the cuboid.

$V = L \times B \times H$ where V is volume, L is length, B is breadth and H is height.

The area of a circle is $A = \pi r^2$ where r is the radius of the circle.

Angles

There are 180° on a straight line.

There are 180° inside a triangle.

An isosceles triangle is a triangle with 2 equal sides and 2 equal angles.

The sum of all the angles inside a n sided polygon is $180(n - 2)$

Pie Chart

In a pie chart, the total angle that corresponds to the entire data set is 360°

Probability

The sum of the probabilities of all outcomes equals 1.

Average Speed

$$\text{Average Speed} = \frac{\text{Distance}}{\text{Time}}$$

Estimate for the mean of a grouped frequency distribution

Estimated mean = sum of (mid interval values multiplied by their frequency) divided by the sum of all the frequencies.

Pythagoras' Theorem

If a , b and c are the sides of a right angled triangle shown below, then

$$a^2 + b^2 = c^2$$

